

Key

2014-2015
ABCALL

HOPKINS

PRETEST

DERIVATIVES

A.P. Calculus ~~Test Two~~

Section One

Multiple-Choice

Calculators Allowed

Time—45 minutes

Number of Questions—15

The scoring for this section is determined by the formula

$$[C - (0.25 \times I)] \times 1.8$$

where C is the number of correct responses and I is the number of incorrect responses. An unanswered question earns zero points. The maximum possible points earned on this section is 27, which represents 50% of the total test score.

Directions: Solve each of the following problems, using the available space for scratch work. After examining the form of the choices, decide which is the best of the choices given and fill in the corresponding choice on your answer sheet. Do not spend too much time on any one problem.

Good Luck!

NAME:

1. $\lim_{x \rightarrow \infty} \frac{5x^2}{3x^2 + 100000x} =$

- A) 0
 B) 0.005
 C) 1
 D) 1.667
 E) does not exist

D

 $\frac{5}{3}$

2. Which of the following functions are not differentiable at $x = \frac{2}{3}$?

I. $f(x) = \sqrt[3]{x-2}$

II. $g(x) = |3x-2|$

III. $h(x) = |9x^2-4|$

- A) I only
 B) II only
 C) I and II only
 D) II and III only
 E) I and III only

D

3. If $y = (\ln x)^3$, then $dy/dx =$

- A) $\frac{3}{x}(\ln x)^2$
 B) $3(\ln x)^2$
 C) $3x(\ln x)^2 + (\ln x)^3$
 D) $3(\ln x + 1)$
 E) None of these

A

$$3(\ln x)^2 \left(\frac{1}{x}\right)$$

4. If $F(x) = x \sin x$, then find $F'(3\pi/2)$.

A) 0

B) 1

C) -1

D) $3\pi/2$ E) $-3\pi/2$

$$x \cos x + \sin x$$

$$\frac{3\pi}{2} \cos \frac{3\pi}{2} + \sin \frac{3\pi}{2}$$

-1

5. $\frac{d}{dx} \left(\frac{1}{x^3} - \frac{1}{x} + x^2 \right)$ at $x = -1$ is

(A) -6

(B) -4

(C) 0

(D) 2

(E) 6

$$\frac{d}{dx} [x^{-3} - x^{-1} + x^2]$$

$$-3x^{-4} + x^{-2} + 2x = -3 + 1 - 2$$

6. The slope of the tangent to the curve $y^3x + y^2x^2 = 6$ at the point $(2, 1)$ is

A) $-\frac{3}{2}$

B) -1

C) $-\frac{5}{14}$ D) $-\frac{3}{14}$

E) 0

$$y^3 + x^3y^2y' + y^2 \cdot 2x + x^2 \cdot 2yy' = 0$$

$$1 + 6y' + 4 + 8y' = 0$$

$$y' = -\frac{5}{14}$$

SCOPE $\frac{f(b)-f(a)}{b-a}$

7. The average rate of change of $f(x) = x^3$ over the interval $[a, b]$ is

- A) $3b + 3a$
- B) $b^2 + ab + a^2$
- C) $\frac{b^2 + a^2}{2}$
- D) $\frac{b^3 - a^3}{2}$
- E) $\frac{b^4 - a^4}{4(b-a)}$

B

Diff. of perfect cubes

$$\frac{b^3 - a^3}{b-a} = \frac{(b-a)(b^2 + ab + a^2)}{(b-a)}$$

8. The function

$$f(x) = \begin{cases} 4 + x^2 & x \leq 1 \\ mx + b & x > 1 \end{cases}$$

is continuous and differentiable for all real numbers. What must be the values of m and b ?

- A) $m = 2, b = 1$
- B) $m = 2, b = 5$
- C) $m = -2, b = 1$
- D) $m = -2, b = 5$
- E) None of these

Cont.

$$3 = m + b \quad -2 = m$$

$$4 = -2 + b$$

$$5 = b$$

9. If $f(x) = -x^2 + x$, then which of the following expressions represents $f'(x)$?

- A) $\lim_{h \rightarrow 0} \frac{-x^2 + x + h - (-x^2 + x)}{h}$
- B) $\lim_{h \rightarrow x} \frac{-x^2 + x + h - (-x^2 + x)}{h}$
- C) $\lim_{h \rightarrow 0} \frac{[-(x+h)^2 + (x+h)] - (-x^2 + x)}{h}$
- D) $\lim_{h \rightarrow 0} \frac{[-(x+h)^2 + (x+h)] - (-x^2 + x)}{h}$
- E) None of these

a

$$\lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

$$f(x) = -x^2 + x$$

$$f(x+h) = -(x+h)^2 + (x+h)$$

10. If $f(x) = \tan(2x)$, then $f'\left(\frac{\pi}{6}\right) =$

$\sec^2(2x) \cdot 2$

$\frac{2}{\cos 2x \cos 2x} = \frac{2}{\frac{1}{2} \cdot \frac{1}{2}}$

- (A) $\sqrt{3}$ (B) $2\sqrt{3}$ (C) 4 (D) $4\sqrt{3}$ (E) 8

11. If $f(x) = (x^2 - 2x - 1)^{\frac{2}{3}}$, then $f'(0)$ is

$\frac{2}{3}(x^2 - 2x - 1)^{-\frac{1}{3}}(2x - 2) = \frac{4(x-1)}{3\sqrt[3]{x^2 - 2x - 1}} = \frac{-4}{-3}$

- (A) $\frac{4}{3}$ (B) 0 (C) $-\frac{2}{3}$ (D) $-\frac{4}{3}$ (E) -2

12. If $f(x) = (x-1)^2 \sin x$, then $f'(0) =$

- (A) -2 (B) -1 (C) 0 (D) 1 (E) 2

$(x-1)^2 \cos x + \sin x (2)(x-1)$

13. If $y = \arctan(\cos x)$, then $\frac{dy}{dx} =$

$\frac{-\sin x}{1 + \cos^2 x} = \frac{-\sin x}{1 + \cos^2 x}$

- (A) $\frac{-\sin x}{1 + \cos^2 x}$ (B) $-(\operatorname{arcsec}(\cos x))^2 \sin x$ (C) $(\operatorname{arcsec}(\cos x))^2$
 (D) $\frac{1}{(\arccos x)^2 + 1}$ (E) $\frac{1}{1 + \cos^2 x}$

14. $\frac{d}{dx}(\arcsin 2x) =$

$\frac{2}{\sqrt{1-4x^2}}$

- (A) $\frac{-1}{2\sqrt{1-4x^2}}$ (B) $\frac{-2}{\sqrt{4x^2-1}}$ (C) $\frac{1}{2\sqrt{1-4x^2}}$
 (D) $\frac{2}{\sqrt{1-4x^2}}$ (E) $\frac{2}{\sqrt{4x^2-1}}$

If $y = \cos^2 3x$, then $\frac{dy}{dx} =$

$2 \cos 3x (-\sin 3x)(3)$

- (A) $-6 \sin 3x \cos 3x$ (B) $-2 \cos 3x$ (C) $2 \cos 3x$
 (D) $6 \cos 3x$ (E) $2 \sin 3x \cos 3x$

16. All the functions below, except one, have the property that $f(x)$ is equal to its fourth derivative, $f^{(4)}(x)$. Which one does not have this property?

- A) ~~$f(x) = \sin x$~~ \cos $-\sin x$ $-\cos x$ $\sin x$
 B) $f(x) = \cos x$
 C) $f(x) = -5e^x$ $-5e^x$ $-5e^x$
 D) $f(x) = e^{2x}$ $e^{2x} \cdot 2$ $2e^{2x} \cdot 2$
 E) $f(x) = e^{-x}$

17. If $g(t) = \frac{\ln t}{e^t}$, then $g'(t) =$

- A) $\frac{1 - \ln t}{e^t}$
 B) $\frac{1 - t \ln t}{e^t}$
 C) $\frac{t \ln t - 1}{te^t}$
 D) $\frac{1 - t \ln t}{te^t}$
 E) $\frac{1 - e^t \ln t}{e^{2t}}$
- $$\frac{e^t \frac{1}{t} - \ln t e^t}{e^t e^t} = \frac{\frac{1}{t} - \ln t}{e^t} = \frac{1 - t \ln t}{te^t}$$

18. If $H(x) = x^3 - x^2 + \frac{1}{x}$, which of the following is $H''(2)$?

- A) $\frac{31}{4}$
 B) $\frac{39}{4}$
 C) $\frac{79}{8}$
 D) $\frac{81}{8}$
 E) $\frac{41}{4}$

$$H' = 3x^2 - 2x - x^{-2}$$

$$H'' = 6x - 2 + \frac{2}{x^3}$$

$$= \frac{10}{1} + \frac{1}{4}$$

$$\frac{41}{4}$$